

## Ronak M. Patel, MD

# Proximal Humerus – Non-Operative Treatment <u>Rehabilitation Guidelines</u>

## **Proximal Humerus Fractures: Nonoperative Treatment**

Outpatient Physiotherapy Phase 1: (Weeks 2-4) ROM

- · Cervical, elbow and wrist ROM
- · Pendulum exercises
- · Instruct in home program Strengthening
- · No cuff strengthening
- · Begin and instruct in program of postural correction
- May begin scapular retraction and depression
- · Grip strengthening Sling
- Arm in sling at all times except for exercises and bathing; Includes sling at night (sleeping in recliner

chair optimal); Discontinue sling between 2-4 weeks

## Outpatient Physiotherapy Phase 2: (Weeks 5-8)

### ROM

- Begin self-assisted forward elevation to 90° and progress in 20° increments per week
- May use pulleys
  - · Begin self-assisted ER with progressive return to full in 20° increments per week
  - · IR in scapular plane as tolerated (No IR behind back)
  - · No cross body adduction
  - · Grade I-II scapulothoracic and glenohumeral mobilizations Strength
  - · No cuff strengthening
  - · Continue scapular retraction and depression
  - · Lower extremity aerobic conditioning Other
  - · Modalities to decrease pain and inflammation
  - · Cryotherapy as necessary

### Outpatient Physiotherapy Phase 3: (Weeks 9-12) ROM

- · Progressive return to full forward elevation and external rotation
- May begin posterior capsular stretching program



550 W Ogden Ave Hinsdale, IL Phone: (630) 323-6116 Fax: (630) 323-6169 For: Test Test DOB: 01/01/1990 MRN: 333245 Page: 2

- May begin IR behind back
- · Grade III-IV glenohumeral and scapulothoracic mobilizations
- · Begin anterior chest wall stretches (pectoralis minor) Strength
- · Instruct in home program and begin submaximal isometrics in flexion, abduction, IR, ER and extension
- · Add progressive isotonics with low resistance, high repetitions as tolerated
- · Emphasize anterior deltoid strength and scapular stabilization
- · Emphasize upper trapezius, serratus anterior force couple rehabilitation to create stable scapular base
- · Assess for and correct compensatory movement patterns
- · UBE with low resistance
- · Continue aerobic conditioning

### Outpatient Physiotherapy Phase 4: (> Week 12) ROM

- · Progressive return to full motion in all planes
- Emphasize posterior capsule stretching
- Maintenance home flexibility program Strength
- · Continue rotator cuff and scapular strengthening program
- Progressive increase in resistance as strength improves
  - Continue UBE with progressive resistance as tolerated
  - · Recreation/vocation specific rehabilitation
  - · Maintenance aerobic conditioning home program