

Discoid Meniscus

The meniscus is a wedge-shaped piece of cartilage that sits in between the bones of the knee and acts as a cushion to protect the bones during movement. Each knee has two menisci.

Healthy menisci are shaped like crescent moons (in fact, "meniscus" comes from the Greek word for crescent). A discoid meniscus is thicker than normal, and often oval or disc-shaped. It is more prone to injury than a normally shaped meniscus.


Some people with discoid meniscus may go through their entire lives and never experience any problems. Many people, however, will have knee problems related to the discoid meniscus. Symptoms often begin during childhood.

Anatomy

The meniscus acts as a "shock absorber" between your femur (thighbone) and tibia (shinbone). It protects the thin articular cartilage that covers the ends of the bones and helps the knee to easily bend and straighten.

There are two menisci in the knee: the medial meniscus on the inside of the knee, and the lateral meniscus on the outside.

The menisci are attached to the tibia bone by a strong tissue called the menisofemoral ligament. This ligament also provides a blood supply to a small portion of the meniscus.



Description

In many cases, a discoid meniscus is shaped like a half moon or complete circle, rather than a crescent moon. It most often occurs on the lateral side (outside) of the knee, and can sometimes be found in both knees.

Types of Discoid Meniscus

There are three types of discoid menisci:

- **Incomplete.** The meniscus is slightly thicker and wider than normal.
- **Complete.** The meniscus completely covers the tibia.
- **Hypermobile Wrisberg.** This occurs when the meniscofemoral ligament that attaches the meniscus to the tibia is absent. Without this ligament, even a fairly normally shaped meniscus can sometimes slip into the joint and cause pain, as well as locking and popping of the knee.

Discoid Meniscus Injuries

A discoid meniscus is more prone to injury than a normal meniscus. The thick, abnormal shape of a discoid meniscus makes it more likely to get stuck in the knee or tear. If the meniscofemoral ligament attachment to the tibia is also missing, the risk for injury is even greater.



Once injured, even a normal meniscus is difficult to heal. This is because the meniscus lacks a strong blood supply and the nutrients that are essential to healing cannot reach the injured tissues.

In many cases of discoid meniscus, patients experience symptoms without there being any injury to the meniscus.

Cause

The cause of discoid meniscus is not known. It is a congenital (present at birth) defect.

Injuries to the discoid meniscus often occur with twisting motions to the knee, such as during sports that require pivoting or fast changes in direction.

Symptoms

The most common symptoms of a discoid meniscus or torn discoid meniscus are:

- Pain
- Stiffness or swelling
- Catching, popping, locking of the knee
- Feeling that the knee is "giving way"
- Inability to fully extend (straighten) the knee

Doctor Examination

Medical History and Physical Examination

After discussing your child's medical history and any events that occurred before symptoms started, your doctor will examine your child's knee.

Your child may or may not have tenderness where the bones meet.

To test for discoid meniscus, your doctor will twist your child's knee with the knee bent and straightened. In many cases of discoid meniscus, there is a popping or clunking sensation. This can sometimes even be heard. In extreme cases, part of the meniscus will pop out of the knee joint and can be seen right under the skin.

Imaging Tests

X-rays. Although x-rays do not show tears of soft tissues like the meniscus, they can show whether there are other problems in the knee. In addition, because the discoid meniscus is so thick, the space between the femur and tibia on the lateral part of the knee may appear widened on an x-ray.

Magnetic resonance imaging (MRI) scan. This study can create detailed images of the soft tissues of the knee joint and is the best imaging test for discoid meniscus.

The MRI can show the abnormal shape of the discoid meniscus, as well as tears within the meniscus. However, the rare Wrisberg type of discoid meniscus cannot be tested with an MRI scan because the problem only occurs when the patient is moving.

An MRI study involves lying still for 30-45 minutes. Many young patients require some sedation or anesthesia to complete the test.

Treatment

Sometimes, a doctor discovers a discoid meniscus when evaluating the knee for a different problem. If the discoid meniscus is not causing any symptoms, then specific treatment for it may not be necessary.

When a discoid meniscus is causing pain, popping, or other symptoms, however, your orthopedic surgeon will probably recommend arthroscopic surgery.

Surgical Treatment

Knee arthroscopy is one of the most commonly performed surgical procedures.

During arthroscopy, the surgeon makes a few small incisions around the knee and inserts a small camera, called an arthroscope, into the joint. The camera displays pictures on a television screen, and the surgeon uses these images to guide miniature surgical instruments.

Most arthroscopic surgeries are done on an outpatient basis. Patients usually go home a few hours after the procedure.

Anesthesia. To prevent pain during the procedure, the patient is given anesthesia. There are different types of anesthesia: Local and regional types of anesthesia numb just parts of the body, and the patient remains awake. General anesthesia puts the patient to sleep. Most children are given general anesthesia for arthroscopic surgery.



Surgical Procedure. Treatment will depend upon the type of discoid meniscus.

- Complete and incomplete discoid menisci with no tears are typically treated with saucerization, a procedure in which the meniscus is cut and re-shaped into a crescent.
- If the discoid meniscus is also torn, the surgeon may perform a saucerization and then trim away the torn portion. Some tears can be repaired with stitches, rather than removed.
- The hypermobile Wrisberg form of discoid meniscus is saucerized if necessary, then stabilized with stitches to sew the meniscus to the lining of the joint.

Rehabilitation

After surgery, your doctor may put your child's knee in a brace or a soft bandage wrap. He or she may need to use crutches for a short period of time. Very young children may need a wheelchair for several weeks because they do not have the balance or strength to use crutches.

Once the initial healing is complete, your doctor may prescribe physical therapy exercises to restore strength and mobility. These may be done at home or with a physical therapist.



Recovery

Most patients return to normal daily activities after arthroscopy for a discoid meniscus. However, if the entire meniscus was removed, there is a risk for continued pain and, potentially, early arthritis.

Surgeons sometimes recommend that patients avoid sports that overstress the knee with cutting type movements. These include soccer, tennis, basketball, and football.

Statistical data on this webpage was reviewed by the AAOS Department of Research and Scientific Affairs.

Last reviewed: October 2014



Reviewed by members of the Pediatric Orthopaedic Society of North America

The Pediatric Orthopaedic Society of North America (POSNA) is a group of board eligible/board certified orthopaedic surgeons who have specialized training in the care of children's musculoskeletal health. One of our goals is to continue to be the authoritative source for patients and families on children's orthopaedic conditions. Our Public Education and Media Relations Committee works with the AAOS to develop, review, and update the pediatric topics within OrthoInfo, so we ensure that patients, families and other healthcare professionals have the latest information and practice guidelines at the click of a link.

AAOS does not endorse any treatments, procedures, products, or physicians referenced herein. This information is provided as an educational service and is not intended to serve as medical advice. Anyone seeking specific orthopaedic advice or assistance should consult his or her orthopaedic surgeon, or locate one in your area through the AAOS "[Find an Orthopaedist](#)" program on this website.

Copyright 2014 American Academy of Orthopaedic Surgeons

Related Articles

Knee Arthroscopy (<http://orthoinfo.aaos.org/topic.cfm?topic=A00299>)

Meniscus Tears (<http://orthoinfo.aaos.org/topic.cfm?topic=A00358>)

OrthoInfo

The American Academy of Orthopaedic Surgeons
9400 West Higgins Road
Rosemont, IL 60018
Phone: 847.823.7186
Email: orthoinfo@aaos.org