

The Thrower's Elbow



Ronak M. Patel, MD^a, T. Sean Lynch, MD^a, Nirav H. Amin, MD^a, Gary Calabrese, PT^a, Stephen M. Gryzlo, MD^b, Mark S. Schickendantz, MD^{a,*}

KEYWORDS

- Overhead throwing athlete • Ulnar collateral ligament • Tommy John • Ulnar subluxation
- Olecranon impingement • Capitellar OCD

KEY POINTS

- Most elbow injuries occur as a result of the stresses incurred during the acceleration phase.
- During overhead throwing, a large valgus force on the elbow created by humeral torque is countered by rapid elbow extension, creating significant tensile stress along the medial compartment, shear stress in the posterior compartment, and compressive stress in the lateral compartment.
- The docking technique in ulnar collateral ligament (UCL) reconstruction demonstrated a lower complication rate and a greater rate of return to play compared with the Jobe technique.
- During surgical management of valgus extension overload syndrome, it is recommended that only the osteophyte and no native olecranon be removed to prevent iatrogenic instability.

INTRODUCTION

The elbow undergoes significant stress during the throwing motion of an overhead athlete. The forces generated in the various phases of the throwing arc are distributed through the soft tissue and bone of the elbow joint. In athletes, such as baseball players, repetition leads to attritional damage to the elbow. The specific constellation of injuries suffered in baseball and other overhead sports, such as softball, football, tennis, and javelin, are well documented. These injuries include (1) medial UCL tears, (2) ulnar neuritis, (3) flexor-pronator injury, (4) medial epicondyle apophysitis or avulsion, (5) valgus extension overload syndrome with olecranon osteophytes, (6) olecranon stress fractures, (7) osteochondritis dissecans (OCD) of the capitellum, and (8) loose bodies.¹

Approximately 55% of high school students participate in sports, and in 2013, softball and baseball ranked as the fourth and third most popular high school sports for girls and boys, respectively.² More than 2 million high school athletes are seen for sports-related injuries on an annual basis. Although the rate of elbow injuries is

relatively low, the total number of such injuries is significant due to the high number of participants. With increasing rates of adolescent and young adults participating in athletics, knowledge regarding the diagnosis and treatment of thrower's elbow remains prudent. The elbow joint is complex, however, and understanding the management of thrower's elbow injuries begins with understanding the anatomy and pathophysiology.

The purpose of this review article is to describe the biomechanics of the throwing motion, the examination of the elbow, the diagnostic evaluation, and the diagnosis and treatment of the spectrum of elbow injuries common to a thrower (**Box 1**).

FUNCTIONAL ANATOMY

The elbow is a ginglymus joint that allows flexion-extension through the ulnohumeral articulation and pronation-supination through the radiocapitellar articulation. It is one of the most congruent joints in the body, with the trochlea covered by articular cartilage over a 300° arc. The bony anatomy of the proximal ulna and olecranon fossa provides primary stability at opposite ends of terminal

^a Sports Health, Department of Orthopaedic Surgery, The Cleveland Clinic Foundation, 5555 Transportation Boulevard, Garfield Heights, OH 44125, USA; ^b Department of Orthopaedic Surgery, Feinberg School of Medicine, Northwestern University, 676 North St. Clair, #1350, Chicago, IL, USA

* Corresponding author.

E-mail address: SCHICKM@ccf.org

Box 1**Thrower's elbow pain differential diagnosis**

UCL injury
 Ulnar neuropathy
 Flexor-pronator injury
 Medial apophysitis or epicondyle avulsion
 Valgus extension overload syndrome
 Olecranon stress fracture
 OCD of the capitellum

motion: less than 20° and greater than 120° of flexion. The radial head provides secondary restraint to valgus stress at 30°. The primary stability during the functional arc of an overhead athlete (20°–120°) emanates from the soft tissue restraints.^{3–5} Furthermore, much of the stability derived from the osseous structure resists against varus stress with the elbow in extension.^{5–7}

The soft tissue structures that provide static valgus elbow stability vital to overhead throwing

include the anterior joint capsule, the UCL complex, and the radial collateral ligament complex. The UCL is composed of 3 bundles: an anterior, a transverse oblique and a posterior. The anterior bundle provides valgus stability throughout the entire range of motion (ROM) and consists of anterior and posterior bands that originate from the inferior aspect of the medial epicondyle and insert at the sublime tubercle on the medial aspect of the coronoid process (Fig. 1A–D).^{5–8} The anterior band provides primary stability against valgus stress from full extension to 90° of flexion and secondary restraint at flexion greater than 90°. The posterior band, which is nearly isometric, provides functionally important restraint between 60° and full flexion, an arc of motion pivotal in the motion of an overhead throwing athlete.^{9,10}

The oblique bundle (transverse ligament) lies at the distal-medial aspect of the joint capsule and does not actually cross the elbow joint. The posterior bundle is thinner and weaker than the anterior bundle and provides secondary elbow stability at greater than 90° of flexion.^{5,6,9}




Fig. 1. Anatomy of the anterior bundle of the medial UCL of the elbow. Illustrations demonstrating traditional anatomy of the medial ulnar collateral ligament anterior band (A), traditional ulnar footprint (dashed line) (B). Illustrations of recently identified anatomy of the medial ulnar collateral ligament anterior band (C) and footprint at an osseous ridge that extends from the sublime tubercle to just medial to the ulnar insertion of the brachialis muscle tendon¹⁶⁰ (outlined with dashed line) (D). (Courtesy of Clinic Center for Medical Art & Photography © 2014, Cleveland. All Rights Reserved; with permission.)

The dynamic elbow stabilizers consist of the muscles in the flexor-pronator mass that originate off the medial epicondyle. This mass consists of the pronator teres, flexor carpi radialis, palmaris longus, flexor digitorum superficialis, and flexor carpi ulnaris (FCU), which functionally stabilize against valgus stress during active motion.¹¹

Lastly, the ulnar nerve courses the medial elbow joint emanating from the arcade of Struthers and passing into the posterior compartment of the arm through the medial intermuscular septum. The nerve, along with a rich plexiform of vessels, enters the cubital tunnel just posterior to the medial epicondyle. The UCL complex forms the floor of the cubital tunnel whereas the roof is composed of the arcuate (Osborne) ligament. Distally, the ulnar nerve passes between the 2 heads of the FCU and rests on the flexor digitorum profundus.

BIOMECHANICS OF THROWING

Overhead throwing sports are typically grouped together because the general motion is similar. Thus, a baseball pitcher's throwing motion, which is the most heavily investigated model, serves as the basis for understanding biomechanics. The baseball pitch is divided into 6 stages of coordinated upper extremity, trunk, and lower extremity movements (Fig. 2).^{7,12–18} The stages specific to elbow motion include

- I. Wind-up: the elbow is flexed and the forearm is pronated as the arm is initially overhead and returns to an adducted position.
- II. Early cocking: the elbow maintains flexion and forearm pronation as the glenohumeral joint is abducted and externally rotated and the leading lower extremity is advanced.

- III. Late cocking: involves increasing elbow flexion between 90° and 120° and forearm pronation to 90° while maximizing shoulder abduction and external rotation.
- IV. Acceleration: the elbow is rapidly extended as the humerus adducts and internally rotates during a coordinated forward movement of the upper extremity and trunk. This stage terminates with ball release over 40 to 50 milliseconds, during which the elbow accelerates as much as 600,000°/s.^{5,19}
- V. Deceleration: rapid deceleration occurs at a rate of 500,000°/s² over a time span of 50 milliseconds as excess kinetic energy is dissipated.^{7,12–16,19}
- VI. Follow-through: the elbow reaches full extension and throwing motion terminates.

Most elbow injuries occur as a result of the stresses incurred during stage IV or the acceleration phase where valgus forces reach as high as 64 Nm.²⁰ The valgus torque concentrate to the medial elbow, primarily the anterior bundle of the UCL.^{11,14} Approximately 300 N of shear force is experienced by the medial elbow.²⁰ Concomitantly, compressive forces at the lateral radiocapitellar joint reach 500 N.²⁰ Nevertheless, modification to throwing biomechanics may not necessarily lead to improved clinical outcomes because the stresses from repetitive throwing may be the driving force to injury.

DEVELOPMENTAL CHANGES OF THE ELBOW

The repetitive stresses at the elbow and shoulder from throwing can lead to developmental changes, and, eventually, injury in young athletes. Small adaptive changes proximally may affect more distal segments of the kinetic chain.²¹ For instance, Garrison and colleagues²² found deficits




Fig. 2. Overhead throwing phases. (Courtesy of Clinic Center for Medical Art & Photography © 2014, Cleveland. All Rights Reserved; with permission.)

in total ROM of the shoulder were associated with UCL tears in a cross-sectional study of high school and collegiate baseball players. Changes in the shoulder include increase in external rotation from humeral retroversion and capsular laxity as well as decrease in internal rotation from osseous adaptations.^{23–26} Polster and colleagues²⁷ demonstrated that the mean dominant arm humeral torsion in professional pitchers was $38.5^\circ \pm 8.9^\circ$ (range, 23.0° – 53.9°) compared with $27.6^\circ \pm 8.0^\circ$ (range, 11.8° – 45.0°) in the nondominant arm. This adaptation may be protective because the investigators found a higher incidence of severe injuries in players with lower degrees of dominant torsion. Burkhart and colleagues²⁸ proposed that increased humeral torsion leads to greater external rotation of the shoulder during late cocking, thus providing a longer throwing arc and potentially a greater peak velocity that increases the stresses experienced by the elbow. Distally at the elbow, Hang and colleagues²⁹ found that 94% of competitive young baseball players had radiographic signs of medial epicondylar apophyseal hypertrophy.

In overhead throwing athletes, understanding the difference between commonly seen asymptomatic adaptive changes and clinically significant pathology is critical in providing proper care to these athletes.³⁰ In a study of asymptomatic professional baseball players, Kooima and colleagues³¹ found an 87% prevalence of chronic UCL injury and an 81% prevalence of posteromedial osteochondral injury. In asymptomatic major league baseball pitchers, increased medial laxity on valgus stress is not uncommon.^{32,33} In a skeletally immature or adolescent thrower, the physis or

apophysis absorbs the stresses of throwing and undergoes changes.^{29,34} With time, asymptomatic changes may progress to symptomatic pathology with increased stress or frequency beyond reparative potential.

PATHOPHYSIOLOGY OF ELBOW INJURIES

King and colleagues³⁵ described a spectrum of elbow injuries in baseball pitchers from medial tension overload to extension overload to lateral compression overload. These injury patterns can be explained by one mechanism: valgus extension overload syndrome.³⁶ During overhead throwing, a large valgus force on the elbow created by humeral torque is countered by rapid elbow extension creating significant tensile stress along the medial compartment, shear stress in the posterior compartment, and compressive stress in the lateral compartment.^{20,30} Repetitive, near-failure tensile stresses create microtrauma and attenuation anterior bundle of the UCL, leading to progressive valgus instability. Continued shear stress and impingement in the posterior compartment lead to olecranon tip osteophytes, loose bodies, and articular damage to the posteromedial trochlea in the continuum of valgus extension overload syndrome (Fig. 3A, B). As the UCL becomes incompetent, the osseous constraints of the posteromedial elbow become important stabilizers during throwing. Subtle laxity in the UCL also leads to stretch of the other medial structures, including the flexor-pronator mass and ulnar nerve. Extrinsic valgus stresses and intrinsic muscular contractions of the flexor-pronator mass lead to tendonitis. Completing the spectrum of thrower's elbow,




Fig. 3. Valgus extension overload syndrome. (A) Posterior view demonstrating valgus force on the elbow. (B) Lateral view demonstrating posterior olecranon osteophytes. (Courtesy of Clinic Center for Medical Art & Photography © 2014, Cleveland. All Rights Reserved; with permission.)

ulnar neuropathy is common given the superficial position of the nerve. The nerve is susceptible to injury from traction, compression, and irritation at the medial aspect of the elbow. In any overhead throwing athlete, UCL attenuation or failure must be ruled out but should not be the only pathology considered.

HISTORY AND PHYSICAL EXAMINATION

A thorough history starts with knowing the patients, their sport, and their level of competition. Asking an athlete specifically about the chief complaint may help delineate between primary (ie, decreased velocity on pitch from UCL attenuation) and secondary processes (ie, pain from posteromedial impingement). Complaints may include pain, decreased motion, mechanical symptoms (clicking, locking, popping, and so forth), instability, and paresthesias as well as throwing-specific symptoms. Changes in accuracy, velocity, stamina, and strength aid in diagnosis and serve as markers to measure improvement. Timing of symptoms may not always be clear; however, if a specific injury or event occurred, it is important to know when, how, and whether there were any antecedent or prodromal symptoms. Any changes in a training or throwing regimen should be noted, including pitch counts, innings pitched, games pitched, and rest between pitching for baseball players.

The timing, onset, and frequency of pain are important to determine.^{20,37} In athletes with valgus instability, approximately 85% experience pain during the late cocking and early acceleration phases of throwing.³⁸

Physical examination starts with inspection of an athlete's posture, arm position, muscle mass, and skin. Any asymmetries compared with contralateral extremity should be detected. Elbow flexion to approximately 70° allows the greatest intracapsular volume and may be an indication of effusion.⁶ Flexion, at a lesser degree, may be secondary to an extension block from posteromedial olecranon osteophytes. With the elbow in extension and forearm in full supination, the carrying angle can be determined. The normal carrying angle is 11° in men and 13° in women.³⁹ Lastly, inspect the skin for any ecchymoses or prior incisions.

Palpate the olecranon, medial and lateral epicondyles, radial head, and soft spot to establish the important landmarks of the elbow. Tenderness on palpation of these landmarks may indicate acute fracture, stress fracture, or tendonitis. In skeletally immature athletes, tenderness may indicate injury to the apophysis or physis. Lateral olecranon tenderness to palpation may indicate a stress fracture whereas proximal medial

tenderness may be related to impingement. Lastly, palpation of the radial head during an arc of passive supination and pronation can help identify osteochondral defects, joint incongruity, and injury to the annular ligament.

Tenderness over the insertions of the various tendons around the elbow can indicate micro-trauma or inflammation. The flexor-pronator mass lies just distal to the medial epicondyle with the arm at 90°. Having patients actively flex the wrist helps identify the tendinous mass, accentuate any pain, and differentiate from UCL pathology. Ranging the elbow from 90° of flexion to approximately 50° to 70° of flexion helps to displace the flexor-pronator mass anterior and exposing the UCL just posterior. Focal swelling and tenderness along the UCL should be concerning. As discussed previously, the UCL has 3 distinct bundles, with the anterior bundle running from the inferior aspect of the medial epicondyle to the medial aspect of the coronoid process.

Directly posterior and distal to the medial epicondyle lies the cubital tunnel, which encloses the ulnar nerve. Palpation of the ulnar nerve from proximal at the arcade of Struthers to distal at the FCU should not elicit any pain. Furthermore, percussion of the nerve should be benign (Tinel sign), but radiating symptoms into the ulnar hand and 2 digits indicate ulnar nerve pathology. The ulnar nerve may be symptomatic, however, even if tenderness is not appreciated. The elbow should be fully extended and then flexed with and without pressure on the nerve proximal to the medial epicondyle. Anterior subluxation of the ulnar nerve can cause local or radiating discomfort.^{40,41}

Stability of the elbow can be assessed with patients in either the supine or seated position. In the supine position, the humerus is stabilized in maximal external rotation and 30° of flexion.^{4,42,43} With the forearm fully pronated and the elbow flexed 20° to 30° to unlock the olecranon from the olecranon fossa, valgus stress is gradually applied to the elbow and opening is assessed.⁴⁴ Less than 1 mm of opening and a firm endpoint should normally be appreciated during the manual valgus stress test. Physiologic laxity may be present, however, and it is more appropriate to compare the stability with the contralateral extremity. Increased opening of the joint space or reproduction of a patient's pain should raise suspicion of injury to the anterior band of the anterior bundle of the UCL.^{7,38,44}

The milking maneuver tests the posterior band of the anterior bundle of the UCL. In this maneuver, the forearm is supinated fully and the elbow is flexed beyond 90° (approximately 120°) and the humerus is at the athlete's side (Fig. 4).⁷ The



Fig. 4. Milking maneuver for evaluation of the UCL. The forearm is supinated fully and the elbow is flexed beyond 90°. The thumb is then pulled laterally by the athlete's contralateral extremity, creating a valgus force on the elbow. Pain, instability, or apprehension is indicative of injury to the UCL.

thumb is then pulled laterally by the examiner or the athlete's contralateral extremity, creating a valgus force on the elbow. Pain, instability, or apprehension is indicative of injury to the UCL.

Lastly, with the patient in the seated position and the forearm supinated, the elbow is slightly flexed. With one hand on the posterior aspect of the distal humerus and the other hand on the volar forearm, the elbow is rapidly extended while applying a valgus stress.³⁶ Pain with this valgus extension overload test indicates impingement of the posteromedial tip of the olecranon on the medial wall of the olecranon fossa.

IMAGING MODALITIES

Standard anteroposterior, lateral, and oblique radiographs are obtained of the elbow. Radiographs may demonstrate calcification of the UCL, osteophytes adjacent to the UCL, olecranon fossa osteophytes, sclerotic OCD lesions, and/or loose bodies. Fluoroscopy is useful in assessing for medial instability by stressing the elbow and comparing with the contralateral extremity. Asymmetry alone, however, may not be enough to diagnose acute injury to the UCL because asymptomatic pitchers have been found to have some laxity in pitching elbow compared with the

contralateral extremity.^{33,45} Nevertheless, greater than 3 mm of opening is concerning for UCL injury and valgus instability.^{5,7}

Conventional radiographs are not sensitive for detecting stress injuries in bone. The sensitivity of initial radiographs is as low as 15% and becomes positive over time in only 50% of patients.^{46,47}

Radionuclide bone scanning is sensitive but less specific for detecting osseous stress injuries, even in their early stages.^{48–52} Radionuclide technetium Tc 99m diphosphonate triple-phase scanning can provide the diagnosis as early as 2 to 8 days after the onset of symptoms.⁵³

CT can help differentiate between stress fractures and other conditions that may show increased uptake on bone scan (Fig. 5). A CT scan is not sensitive, however, in detecting stress injuries in their early stages.⁵⁴

MRI can detect early stress changes as well as muscle and tendon changes, loose bodies, osteochondral injuries, olecranon osteophytes, and neurologic changes to thrower's elbow. MRI is useful in evaluating UCL avulsions, partial ligamentous injuries, midsubstance tears, and the status of the surrounding soft tissues (Fig. 6). MRI has been found 57% sensitive and 100% specific in detecting UCL injuries.^{55–58} MRI arthrography seems to improve the sensitivity of detection of UCL tears, with saline injection improving the




Fig. 5. Axial slice of a CT scan of a professional pitcher demonstrating a stress fracture of the olecranon (arrow).




Fig. 6. Coronal slice of an MRI demonstrating a medial UCL tear (circle).

sensitivity of UCL detection to 92%.⁵⁶ Potter⁵⁸ and Gaary and colleagues⁵⁷ have reported similar sensitivity and specificity with nonarthrogram MRI using special sequences at the Hospital for Special Surgery. Timmerman and colleagues⁵⁹ compared CT arthrogram with both contrast-enhanced MRI and nonenhanced MRI and found a sensitivity of 86% and specificity of 91%.

ULNAR COLLATERAL LIGAMENT INJURIES

Depending on the extent of damage to the UCL, specific treatment programs can be implemented. Complete disruption of the anterior bundle of the UCL can destabilize the elbow against valgus stress encountered during the throwing motion. Partial tears of the UCL can be managed nonoperatively in low-demand patients^{3,60}; however, results in overhead throwing athletes have not been promising.⁶¹ Overall, treatment options for UCL injury include nonoperative rehabilitation, direct ligament repair, or free-tendon graft reconstruction.

Nonoperative Treatment

After a complete evaluation and diagnosis of a UCL injury through physical examination and radiologic studies, physician and athlete must agree on the appropriate course of care. Nonsurgical treatment measures are indicated for the initial treatment of sprains of the medial UCL in the vast majority of cases. Patients who present with findings consistent with a partial tear of the UCL, grade I and some grade II, should be initially

placed on a period of active rest for 6 to 12 weeks. It is important to protect the elbow from valgus stress, including throwing, for a minimum of 6 weeks. Using a criteria-based rehabilitation program assures that a patient's progress is appropriate for individual rehabilitation potential at each criteria stage.

Initially, athletes are treated with cryotherapy, pain-modulating electrotherapy modalities, antiinflammatory medication, and a hinged elbow brace restricting full extension because relief of pain and reduced inflammation dictate the subsequent rehabilitation strategies. The early focus of rehabilitation is in regaining or maintaining elbow and shoulder ROM in conjunction with shoulder-strengthening exercises. Scapular-based exercises are initiated immediately for both nonoperative and operative UCL rehabilitation programs. Patients can continue core- and lower quarter-strengthening exercises performed without gripping heavy weight or resistance. Once a patient has regained full pain-free elbow ROM, there is a progression from isometric to isotonic upper arm-based to a forearm-based resistance program focusing on strengthening the medial dynamic stabilizers with emphasis on the pronator teres, FCU, and flexor digitorum superficialis. Hamilton and colleagues¹³ found that when UCL stabilizing capabilities are compromised, activity of these medially based muscles is decreased. A criteria-based return-to-throw program is initiated when functional patterns, with resistance, consistent for pitching are pain-free and valgus stress testing is negative. Retting and colleagues⁶² evaluated 31 throwing athletes treated nonoperatively for a UCL injury with a minimum of 3 months of rest with rehabilitation. They reported a 42% return to competitive throwing at the same level or higher at an average of 24.5 weeks.

Recent interest in platelet-rich plasma (PRP) has led to a broad array of applications. Podesta and colleagues⁶³ evaluated their outcomes in 34 athletes with partial UCL tears who received 1 PRP injection to the elbow after failing 2 months of typical nonoperative management; 30 of the 34 returned to play at preinjury level at an average of 12 weeks.

Operative Treatment

Overhead throwing athletes with complete disruption of the anterior bundle of the UCL are candidates for surgical intervention if they wish to return to preinjury level of play. Athletes with partial tears unable to return to competitive throwing (or other overhead sport) due to continued medial elbow pain despite completion of an adequate course of nonoperative treatment are considered

candidates for surgical treatment as well. The goal of surgical reconstruction of the medial UCL is restoration of valgus stability to the elbow.

Direct primary repair of the UCL is reserved for acute avulsion injuries from either the humeral origin or coronoid insertion.^{38,64,65} UCL injuries in throwing athletes are typically present, however, as attenuated ligaments or midsubstance tears. Chronic repetitive microtrauma leads to significant scarring and subsequent inability for effective primary repair of the UCL.^{61,66} Limited studies of results after primary repair of medial UCL injuries exist.^{38,67,68} Level IV retrospective series have shown inconsistency in documentation of athletic level and return to play.^{65,66} A case series of 47 adolescent athletes (mean age 17.2 years) by Savoie and colleagues⁶⁸ reported 93% good to excellent results after primary repair of proximal and distal ligament avulsion injuries using suture anchors or bone tunnels.

Meanwhile, studies comparing repair with reconstructive techniques have shown better results with the latter. Conway and colleagues³⁸ treated 14 overhead throwing athletes with UCL deficiency with primary repair and 56 with graft reconstruction. In the repaired group, 50% of the athletes returned to preinjury level of sport, and, overall, 71% had good or excellent results. In the reconstruction group, 68% returned to preinjury level of sport, whereas 80% had good or excellent results. Andrews and Timmerman⁶⁹ evaluated 72 professional baseball players who underwent elbow surgery and found that neither of the 2 athletes who underwent primary repair of the UCL returned to sport, whereas 12 of the 14 who underwent reconstruction were able to return to play. Lastly, Azar and colleagues⁴⁵ reported results of 67 patients treated with UCL primary repair or reconstruction with 12- to 72-month follow-up; 5 of the 8 patients (63%) treated with repair returned to preinjury level of play compared with 48 of 59 (81%) in the reconstruction group.

Frank Jobe first performed reconstruction of the elbow medial UCL on September 25, 1974, on Los Angeles Dodgers left-handed pitcher Tommy John. John returned to pitching in 1976 and over the next 13 seasons went on to pitch 2500 innings, compiling a record of 164 wins and 125 losses, and never having another significant problem with his elbow. John's successful return to pitching after surgery revolutionized the treatment of athletes with injuries to the medial UCL and popularized the procedure, known as *Tommy John surgery*.

In the Jobe 3-ply technique, the ipsilateral palmaris longus tendon is harvested as a graft.^{38,64} Other suitable options for graft material include the contralateral palmaris longus tendon and

gracilis tendon. Allograft gracilis tendon may also be used. Savoie and colleagues⁷⁰ performed hamstring allograft medial UCL reconstruction in 116 overhead athletes. Of these 116 athletes, 110 returned to play with 88% playing at or above preinjury level.

Two converging drill holes are created in the sublime tubercle of the proximal ulna, creating a bone tunnel, and 2 divergent drill holes are created in the medial epicondyle (**Fig. 7**). With the assistance of a suture passer, the graft is passed first through the bone tunnel in the ulna. It is then crossed over itself as the posterior limb of the graft is passed up the anterior tunnel into the medial epicondyle with the anterior limb going into the posterior tunnel. The limb within the posterior tunnel is then brought up around the back of the epicondyle and passed distally back into the anterior tunnel, exiting at the entrance point into the medial epicondyle. With the elbow positioned in approximately 30° of flexion and the forearm in neutral rotation, with a slight varus moment applied to the elbow, the graft is tensioned and sutured to itself, resulting in a 3-ply graft reconstruction. Any remaining native ligament is then incorporated into the graft, reinforcing the construct.

The classic Jobe surgical technique involves exposing the UCL by reflecting the flexor-pronator origin off the medial epicondyle and anteriorly transposing the ulnar nerve in a submuscular position. Early published results of medial UCL reconstructions performed using this technique reported a high incidence of postoperative ulnar nerve complications.^{38,64} Thompson and colleagues reported a 31% incidence of postoperative ulnar nerve dysfunction whereas Smith and colleagues found an incidence of 21%. In an effort to minimize these complications, contemporary techniques use a muscle-splitting approach through the FCU.^{71,72} Additionally, most surgeons currently reserve ulnar nerve transposition for those athletes with clinically significant ulnar nerve instability.⁷²

Several modifications of the Jobe original 3-ply technique have been developed over the past decade.⁷³ In addition, a variety of graft fixation methods have been investigated in both the clinical and laboratory settings. These include interference screws, suture anchors, flip-buttons, and combinations of these fixation methods, along with variations in tunnel placement both proximally and distally.⁷³⁻⁸⁷ An arthroscopically assisted technique has been studied in the laboratory and has shown biomechanically promising results.⁸⁵

The most widely applied and studied of these newer techniques is the docking procedure.^{74,76,77,83,84,86,88,89} In the docking technique,




Fig. 7. Illustration of classic 3-ply medial UCL reconstruction technique as described by Jobe. Inset: Completed reconstruction. (Courtesy of Clinic Center for Medical Art & Photography © 2014, Cleveland. All Rights Reserved; with permission.)

converging drill holes are used to create a tunnel at the level of the sublime tubercle of the proximal ulna as in the Jobe 3-ply method. Instead of divergent tunnels in the medial epicondyle of the humerus, however, a single blind-ended tunnel (socket) is created (Fig. 8). At the end of this socket, 2 small exit holes are created to allow for passage of sutures that have been sutured to end of the graft. These sutures are used to pull the graft into the humeral socket where it is seated at the base. The passing sutures are then tied to each other over the back of the epicondyle. This method of fixation eliminates the suture/graft interface that is present in the 3-ply reconstruction and has been shown in biomechanical studies to have higher peak load to failure values compared with Jobe and interference screw techniques.^{74,83} A recent systematic review by Watson and colleagues⁹⁰ found that the docking technique and a suspensory button technique most commonly failed secondary to suture failure, whereas the most common modes of failure for the Jobe technique and interference screw technique were ulnar tunnel fractures and graft ruptures, respectively.

A modification of the docking technique has recently been introduced. The DANE TJ UCL reconstruction uses traditional docking fixation proximally into a socket within the medial epicondyle of the humerus.^{82,91} Distally, however, instead of a tunnel in the ulna, another socket is created

(Fig. 9). Fixation of the ulnar end of the graft is achieved with an interference screw, whereas fixation proximally is performed with 2 sutures exiting the end of the humeral socket and tied to each other, as in the traditional docking technique. To augment proximal fixation of the graft, some surgeons place an interference screw within the medial epicondyle socket. The addition of an interference screw for fixation proximally has been shown to result in less gap formation under valgus loading in the laboratory setting, perhaps allowing for better healing of the graft within the humeral bone socket.^{78–80} Aperture fixation, as is achieved with interference screws, may also result in increased graft isometry and increased stiffness of the overall construct.⁷⁵ Concerns with this type of fixation include low initial fixation strength and the potential for graft slippage in the early postoperative period.⁷⁸

Surgical reconstruction of the UCL dictates a reduced pace of rehabilitation and is a lengthy process. Therefore, the postoperative management varies depending on the surgical procedure performed. The initial goals of the program center on protection of the UCL graft, decreasing pain and effusion, and maintaining muscular strength in the forearm-based musculature.

Patients are fit postoperatively with a posteriorly based elbow splint fixed at 90° of flexion for immobilization and a compression dressing during the




Fig. 8. Schematic of original docking technique for UCL reconstruction. Inset: Completed reconstruction. (Courtesy of Clinic Center for Medical Art & Photography © 2014, Cleveland. All Rights Reserved; with permission.)

acute phase of healing. During the immediate postoperative, acute phase of healing, the athlete is started on a program emphasizing wrist flexion and extension active ROM gripping exercises,

while maintaining a neutral wrist position, and sub-maximal isometric exercises for the hand, wrist, and elbow in all directions. The initiation of the subacute phase of rehabilitation, usually 2 to




Fig. 9. Drawing of the DANE TJ modification of the docking technique. Inset: Completed reconstruction. (Courtesy of Clinic Center for Medical Art & Photography © 2014, Cleveland. All Rights Reserved; with permission.)

4 weeks, occurs as patients are transferred from the posterior splint to a hinged elbow brace that allows between 40° and 100° of elbow motion. The ROM is gradually increased so that full elbow ROM is achieved by postoperative week 5 to 6, at which time the hinged elbow brace usage is discontinued. Isometric exercise progression is advanced to include moderately applied force and light resistance isotonic exercises. Initiating the isotonic exercises while maintaining decreased valgus load allows for maximum support of the elbow as a new exercise procedure is introduced into the rehabilitation program. The athlete is advanced in the scapular stabilization and shoulder exercise program to include light to moderate resistance below 90° of shoulder elevation. Manual resistance is used for scapular stabilization exercises and short arc proprioceptive neuromuscular facilitation upper extremity patterns are initiated with resistance placed proximal to the elbow.

During the intermediate phase of rehabilitation, 6 to 10 weeks, the exercise program is advanced to include shoulder positioning into external rotation coupled with scapular retraction, shoulder elevation greater than 90°, biceps and triceps isotonic resistance, wrist pronation and supination, and core stabilization. Eccentric loading is initiated manually between 9 and 10 weeks postoperatively. Emphasis is placed on exercises that activate the FCU and flexor digitorum superficialis because they are believed to assist the UCL in medial elbow stabilization. Plyometric exercises are initiated after successful performance of manual and active resisted eccentric loading exercises. The athlete couples the exercise progression with nonthrowing baseball pitching-specific drills for balance point position, arm path to shoulder/elbow 90/90 positioning, and stride direction and length. The shoulder rotator cuff and scapulothoracic exercises incorporate functional chain positioning of the lower extremity and trunk to maximize complex movement patterns necessary for a successful return to throwing. A return-to-throw progression and interval throwing program is initiated at week 16.

Outcomes after UCL reconstruction have shown generally favorable results, with reported return-to-play rates as high as 95%.^{38,45,64,67,72,77,84,88,89,91,92} A systematic review of the literature by Vitale and Ahmad⁹³ concluded that overall 83% of patients (493) had excellent results (return to the same level of play for at least 1 year). An overall complication rate of 10% is reported, with ulnar nerve complications the most common. Better results were seen with a muscle-splitting surgical approach, no ulnar nerve

transposition, and utilization of the docking technique. Also, adolescent athletes do not fare as well as college and professional athletes after surgical reconstruction of the UCL. In a retrospective study of 27 high school athletes who had undergone UCL reconstruction, Petty and colleagues⁹⁴ reported that only 74% were able to return to the same level of play after surgery. They identified grossly positive stress radiographs, sublime tubercle avulsion fractures, and ossicles within the proximal end of the ligament as poor prognostic indicators.

In a more recent systematic review of more than 1300 patients, Watson and colleagues⁹⁰ found the docking technique demonstrated a lower complication rate (6.0%) compared with the Jobe technique (51.4%) ($P = 4.48 \times 10^{-6}$). Additionally, there was a trend toward a greater rate of return to play with the docking technique compared with the Jobe technique at 90.4% and 66.7%, respectively ($P = 1.29 \times 10^{-5}$).

Revision surgery for the treatment of failed reconstructed ligaments is technically challenging and is associated with a high incidence of complications and generally poor outcomes.^{95,96} In a recent case series, Dines and colleagues⁹⁶ reported that only 33% of athletes were able to return to play after revision medial UCL reconstruction, with 40% of patients experiencing complications.

ULNAR NEUROPATHY

Ulnar neuropathy at the elbow is the second most prominent neuropathy of the upper extremity, and its superficial location makes it particularly susceptible to injury in throwing athletes. The ulnar nerve is susceptible to several mechanical factors, including compression, traction, and irritation of the nerve.⁴⁴ During the acceleration phase of the throwing motion, the ulnar nerve is subject to longitudinal traction.⁹⁷ Potential sites of compression proximal to the cubital tunnel include the arcade of Struthers and the medial intermuscular septum. Distal sites of potential compression include the area between the FCU and the medial forearm musculature. Recently, Li and colleagues⁹⁸ even reported compression through the anconeus epitrochlearis, which may be hypertrophied in overhead throwing athletes. Aoki and colleagues⁹⁹ found a hypertrophic medial head of the triceps impinging on the ulnar nerve as the elbow was flexed greater than 90° in a series of adolescent baseball players with ulnar neuropathy. Osborne ligaments represent the predominant compression site in the cubital tunnel, which is bordered laterally by the elbow, anteriorly by the medial epicondyle,

and medially by the origin of the FCU,^{100,101} although compression may also occur from osteophytes, loose bodies, or synovitis.

Repetitive throwing motions can cause both physiologic and pathologic responses that may be the primary cause of ulnar neuropathy.^{102,103} Many secondary causes exist. More than 40% of athletes with valgus instability experience ulnar neuritis secondary to irritation from the inflammation of the UCL as well as increased stretch from valgus stress. Approximately 60% of athletes with medial epicondylitis develop ulnar nerve symptoms.^{38,44,104} Osteophytes from valgus extension overload and friction from ulnar nerve subluxation can serve as underlying etiologies as well.¹⁰⁵

Intraneural pressure within the ulnar nerve varies with wrist, elbow, and shoulder position. Pechan and Julis¹⁰⁶ found the pressure within the nerve to be 3 times the resting level when the elbow was flexed and the wrist extended. Continuation of the throwing motion with further elbow flexion and shoulder abduction causes the intraneural pressure to rise 6 times the resting level. This increased pressure is attributed to nerve stretch, tightening of the cubital tunnel, and compression.¹⁰⁷ Repetitive motions can induce chronic changes to the ulnar nerve and surrounding soft tissues, potentially leading to nerve fibrosis and ischemia.

In addition to the history and examination discussed previously, the elbow flexion test with the elbow maintained in maximum flexion and wrist in extension for 1 minute should be conducted. Symptoms of ulnar neuritis consist of an aching pain along the ulnar side of the forearm radiating into the ulnar 2 digits of the hand. Reports of clumsiness or numbness should be evaluated further. Monofilament testing can detect early sensory changes, and hand intrinsic atrophy or weakness represents the earliest motor changes.

Electrodiagnostic testing may be useful in diagnosing ulnar neuropathy; however, changes may not be seen until disease has advanced. Wei and colleagues¹⁰⁸ performed nerve conduction velocity testing in baseball pitchers and did not find a significant difference between the dominant and nondominant arms among injured pitchers. A negative electrodiagnostic test, however, does not rule out ulnar neuritis.^{41,102}

Nonoperative Treatment

Nonoperative management typically begins with activity restriction, antiinflammatories, cryotherapy, and physical therapy. Ulnar subluxation or dislocation may require a brief period of

immobilization. After resolution of symptoms, a gradual return-to-throw program can be initiated. Symptoms from superficial irritation of the nerve can be treated with elbow pads. Ulnar neuropathy in overhead throwing athletes typically stems, however, from an underlying cause, and as throwing is resumed, symptoms likely resurface.

Operative Treatment

Indications for operative management include failed nonoperative treatment, persistent ulnar nerve subluxation, symptomatic tension neuropraxia, and concomitant medial elbow problems that require surgery.¹⁰⁷ The surgical options for treatment include decompression (in situ) of the ulnar nerve, medial epicondilectomy, and anterior transposition of the nerve (submuscular, intramuscular, or subcutaneous). Simple decompression and medial epicondilectomy are prone to failure in overhead throwing athletes. Decompression may address some of the compressive forces within and around the cubital tunnel but does not address irritation from traction and repetitive motion. Furthermore, in cases of ulnar nerve subluxation, decompression fails to stabilize the nerve. Medial epicondilectomy aims to prevent traction irritation to the nerve as it passes posterior to the medial epicondyle. Well-recognized complications include, however, nerve instability, valgus instability from iatrogenic injury to the UCL, and weakness from disruption of the flexor-pronator mass.

Anterior transposition of the ulnar nerve is the mainstay of surgical management of ulnar neuritis. Subcutaneous transposition has the advantage of less surgical morbidity to the flexor-pronator mass and may be recommended in patients undergoing concomitant UCL reconstruction.^{45,109,110} The main disadvantage of the subcutaneous technique is its susceptibility to direct trauma.^{44,102,103} Other concerns include potential for developing instability and recurrence of symptoms from new compression under the subcutaneous fasciadermal sling. Anterior submuscular transposition involves greater surgical dissection in the medial soft tissues and a lengthier postoperative rehabilitation course but provides thorough decompression of the nerve while protecting it from direct and indirect trauma. In submuscular transfer, the nerve sits deep to the medial flexor mass and arises superficial to the pronator muscle mass. The outcomes of ulnar nerve transposition in overhead athletes support both approaches in throwing athletes. Studies comparing the results of these techniques, including 2 meta-analyses, have not found a significant difference between the procedures in terms of postoperative clinical

outcomes or motor nerve conduction velocities; however, these were not specific to throwing athletes.^{111,112}

Del Pizzo and colleagues⁴¹ reported their findings on 19 baseball players who underwent anterior subcutaneous ulnar nerve transposition for recalcitrant ulnar neuritis; 9 of 15 athletes (60%) evaluated at 3 to 58 months postoperatively were able to return to play. Rettig and Ebben¹¹⁰ performed 21 anterior subcutaneous transposition procedures in 20 athletes for failed conservative management of cubital tunnel syndrome. All athletes returned to play at an average of 12.6 weeks. The investigators recommended subcutaneous transposition in light of faster postoperative recovery and rehabilitation.

The senior authors' (SMG and MSS) preference is subcutaneous anterior transposition of the ulnar nerve supported by facial slings, paying particular care to preserve the motor branches to the FCU. A 4- to 6-cm curvilinear incision is made just posterior to the medial epicondyle along the path of the ulnar nerve. Dissection is carried out, taking care to protect any branches of the medial antebrachial cutaneous nerve. The ulnar nerve is identified in the cubital tunnel by opening the cubital tunnel at its midsection just posterior to the medial epicondyle. The distal one-half of the cubital tunnel is then released, including the Osborne ligament and any compressive fibers in the proximal FCU. The proximal cubital tunnel is then released, extending proximally to the arcade of Struthers. Two fascial slings are developed from the superficial flexor-pronator muscle fascia, and the fascia is closed. The ulnar nerve is transposed anterior to the medial epicondyle and held loosely in place by suturing the fascial slings to the flexor fascia. The posterior triceps fascia is then sutured to the medial epicondyle to close the cubital tunnel, and the FCU fascial split is reapproximated loosely. The elbow is splinted in 90° of elbow flexion for the 7 to 10 days postoperatively to allow soft tissue healing, followed by progressive motion and rehabilitation to return to play.

FLEXOR-PRONATOR INJURY

The flexor-pronator muscle mass at the medial side of the elbow provides dynamic stability against valgus forces.^{13–16} Repetitive contraction of the flexor-pronator muscles occurs during the acceleration phase of throwing as well as with wrist flexion during ball release.¹⁶ An acute complete rupture of the common flexor-pronator origin from the medial epicondyle is an uncommon injury in overhead athletes; rather, athletes may develop a spectrum of injuries from mild muscular overuse

to chronic tendinitis or acute partial muscle tears (Fig. 10).⁶⁵ The typical presentation includes an athlete with medial elbow pain during the late cocking or acceleration phase of throwing with continuation during ball release as the forearm is pronated and the wrist flexed. With complete ruptures of the flexor-pronator muscle mass, players may recall general prodromal symptoms followed by a single event leading to a pop sensation or sound.

The main differential diagnosis in these cases should include injury to the UCL. In flexor-pronator injuries, athletes have tenderness to palpation just distal to the medial epicondyle along the common origin. Meanwhile, in UCL injuries, the tenderness is typically posterior and distal to that of flexor-pronator pain along the sublime tubercle. Lastly, pain with flexor-pronator injury should be exacerbated with wrist flexion and elbow extension. Nevertheless, clinical differentiation between these 2 entities can be difficult and an MRI aids in diagnosis. Combined UCL and flexor-pronator injuries are not uncommon.

Overuse tendonitis and partial tears of the flexor-pronator muscle mass can be treated with active rest, ice, antiinflammatory medications, physical therapy, and gradual return to throwing. Corticosteroids are generally avoided given the proximity of the UCL. No data are currently available on the efficacy of PRP injections for flexor-pronator injury in overhead throwing athletes. Complete ruptures of the flexor-pronator origin require prolonged rest and rehabilitation. Splinting the wrist in neutral position may alleviate acute



Fig. 10. Coronal slice of an MRI of a minor league reliever demonstrating a tear of the flexor-pronator mass near the proximal origin (circle).

pain during the first 7 to 10 days after injury. Rehabilitation with emphasis on ROM first followed by resistance training is necessary before interval throwing is initiated. If pain or weakness reoccurs with throwing, then surgical repair may be necessary. Because surgical treatment is rarely required, failure of nonoperative treatment should heighten awareness of other underlying pathology.

Overhead throwers are also vulnerable to pronator syndrome, which is compression of the median nerve caused by hypertrophy of pronator teres secondary to repetitive activity. Athletes complain of vague, fatigue-like pain over the proximal volar aspect of forearm exacerbated by resisted forearm pronation and wrist flexion. If nonoperative management with activity modification and anti-inflammatory medications fails, then surgical exploration with division of superficial head of the pronator teres and decompression of the median nerve may be required. Similarly, hypertrophy of the flexor-pronator mass can cause a localized compartment syndrome induced by repetitive throwing. Medial elbow and forearm pain is associated specifically with throwing and resolves with rest. This condition may be difficult to diagnose but typically responds well to adequate stretching, interval rests, and adherence to proper pitching mechanics.³

MEDIAL EPICONDYLE AVULSION OR APOPHYSITIS

Little League elbow is a general term referring to medial-sided stress injuries that can occur in skeletally immature throwing athletes.^{113,114} Medial epicondyle avulsion injury and apophysitis are the most common injuries and are prevalent in youth baseball. Hang and colleagues²⁹ found that 52% of Little League players in Taiwan reported medial elbow pain or soreness at some point during the course of a season. Grana and Rashkin¹¹⁵ reported that 58% of older adolescent pitchers experience elbow pain or injury at some point during the season. In skeletally immature throwers, both the static (UCL) and dynamic (flexor-pronator muscle group) medial stabilizers attach to the medial epicondyle, thus conferring all of the static and dynamic stress to the medial epicondylar physis. Valgus and contractile forces across the elbow that result in ligament and tendon injury in the adult thrower lead to injury to the weaker medial epicondyle apophyseal plate in the skeletally immature thrower. Symptomatic injury is associated with the combination of repetitive forces on the medial elbow and inadequate intervals of rest.¹¹⁶⁻¹¹⁸ The use of breaking pitches in this age group and improper throwing

mechanics have also been suggested as possible causes but not proved.¹¹⁷⁻¹¹⁹ Bennett coined the term, Little Leaguer's elbow, to describe the clinical and radiographic changes associated with these medial-sided traction injuries in skeletally immature athletes.¹²⁰ Repetitive valgus loading can lead to apophyseal fragmentation or avulsion of the medial epicondyle apophysis as well as changes in the radiocapitellar joint laterally.

Medial epicondylar apophysitis should be suspected in young athletes with point tenderness over the medial epicondyle and pain with valgus stress of the elbow. Patients with avulsion fractures typically have significant swelling and decreased ROM. Radiographs may show a subtle widening of the physeal plate and comparison views of the uninvolved limb are vital. MRI typically shows more findings than radiographs; however, these do not necessarily have clinical correlation and do not change management.¹²¹

Initial treatment consists of an extended period of rest and cryotherapy until symptoms resolve and a gradual return to throwing. Harada and colleagues¹²² found that athletes noncompliant with pitching restrictions and who returned to rigorous activity prior to bone union on radiographs had significant delays in bone union at both 6 months and 1 year compared with athletes who waited until complete bone union prior to resuming rigorous throwing. Although activity modification is successful in apophysitis, treatments of epicondylar avulsions remain controversial.¹²³

In cases of complete avulsion of the medial epicondyle, most investigators recommend open reduction and internal fixation if the fragment is displaced 5 mm or greater. Nonoperative management is successful in the management of minimally displaced fractures without instability. These injuries can be treated with splint immobilization for 5 to 7 days, followed by early motion.¹²⁴⁻¹²⁶ Lawrence and colleagues¹²⁷ recommend nonoperative management in young athletes with low-energy medial epicondyle avulsions, a stable elbow, and minimal fracture displacement (5.3 ± 2.0 mm). Open reduction and internal fixation can be successful in athletes who sustain more significant trauma, who have elbow laxity or instability, or who have significant fracture fragment displacement (7.1 ± 2.9 mm) after a fracture of the medial epicondyle.

VALGUS EXTENSION OVERLOAD SYNDROME

Valgus extension overload can occur with an attenuated UCL or in a physiologic lax elbow with repetitive valgus stress from throwing. Athletes most commonly complain of posteromedial

elbow pain during the extension (late acceleration) or follow-through phase of throwing.¹²⁸ During these phases, the elbow subluxates and increases force in the lateral and posterior compartments. Continued compressive and rotatory forces in the lateral compartment lead to synovitis and osteochondrosis of the radiocapitellar joint.^{3,36} Posterior and posteromedial olecranon osteophytes that form from impingement may fracture and contribute to loose bodies along with osteochondral fragments from the radiocapitellar joint (Fig. 11). When athletes complain of locking or catching, loose bodies should be considered. Radiographs may demonstrate osteophytes or loose bodies. Cohen and colleagues¹²⁹ have found a typical pattern of valgus extension overload syndrome on MRI findings. All 9 throwing athletes had pathology at the articular surfaces of the posterior trochlea and the anterior, medial olecranon. The investigators also found that these MRI findings correlated with arthroscopic findings.

Nonoperative Treatment

Initial treatment should consist of active rest with cryotherapy, iontophoresis, and antiinflammatory medications to relieve pain. As symptoms subside and motion returns to baseline, rehabilitation should include dynamic stabilization and strengthening exercises with emphasis on eccentric strengthening of the elbow flexors to control rapid extension of the elbow. Manual resistance exercises of concentric and eccentric elbow flexion are performed prior to an interval throwing program.¹³⁰




Fig. 11. Sagittal cut of an MRI of an overhead throwing athlete demonstrating multiple irregularities of the posterior articular surface of the olecranon (square).

Operative Treatment

Andrews and Timmerman⁶⁹ reported posteromedial olecranon impingement the most common diagnosis requiring surgery in baseball players (78%). Failure of nonoperative management for impingement-related pain is a surgical indication. Nonoperative treatment is deferred in overhead athletes with symptomatic posterior medial osteophytes or loose bodies. The treatment of choice in these instances is débridement and removal of the loose bodies. Arthroscopic treatment has become the mainstay of olecranon débridement and excision of loose bodies.^{36,131} Arthroscopic evaluation also allows for débridement or drilling of osteochondral defects, resection of hypertrophic synovium, and inspection for undersurface tears of the UCL.¹³² Athletes must understand the risk of recurrence with continued throwing as well as risk of damage to the ulnar nerve with this technique.⁶⁹

Reddy and colleagues¹³³ reported 187 arthroscopies done in 172 patients at the Kerlan-Jobe Orthopaedic Clinic. The most common diagnosis was posterior impingement (51%), followed by loose bodies (31%) and degenerative joint disease (22%). Laxity of the UCL was seen in 6% of cases. Although 68 patients were lost to follow-up, they reported 49% excellent, 36% good, 11% average, and 4% poor results, based on the modified Figgie score; 47 of the 55 professional athletes (85%) returned to their previous level of competition. There were 3 transient complications, 1 related to the ulnar nerve. Andrews and Timmerman⁶⁹ evaluated 56 professional baseball players who underwent arthroscopic olecranon osteophyte excision either as an isolated procedure or with concomitant ulnar nerve transposition or UCL reconstruction; 23 of 34 patients (68%) available for minimum 24-month follow-up returned to play at least 1 season. However, 14 (41%) required reoperation, including repeat débridement of olecranon osteophytes (6) and UCL reconstruction (5). The investigators cautioned against excessive olecranon excision. Resection of more than 3 mm of the posteromedial olecranon jeopardizes the function of the anterior bundle of the UCL because it exposes a potentially attenuated UCL to higher stresses.^{134,135} Thus, it is recommended that only the osteophyte and no native olecranon be removed.

OLECRANON STRESS FRACTURE

Proximal olecranon stress fractures occur from the repetitive microtrauma, excessive tensile stress from the triceps tendon, and posterior impingement of the olecranon against the olecranon fossa

associated with competitive overhead throwing.¹³⁶ Stress fractures can have posterolateral or posteromedial olecranon pain and tenderness during and after throwing.¹³⁷ There is typically no pain at rest and there is a gradual onset rather than a single event. Schickendantz and colleagues¹³⁷ state pain with percussion of the proximal ulna may indicate a stress fracture because it was a positive test in all 7 athletes with MRI-confirmed stress reactions. Stress fractures of the proximal olecranon should be differentiated from avulsion fractures of the tip of the olecranon and from a persistent olecranon apophysis, which may be treated differently. Standard radiographs as well as advanced imaging may be needed for diagnosis.

Nonoperative Treatment

Initial treatment includes rest, immobilization, and throwing cessation. Specifically, any valgus stress should be avoided for a minimum of 6 weeks. Full extension should also be avoided with the use of a splint or orthosis set to approximately 20° of extension for the first 4 weeks. At 4 weeks, full ROM is allowed and progressive resistance exercises of the elbow are initiated. At 6 weeks, sport-specific rehabilitation is initiated and an interval throwing program typically starts at approximately 8 weeks. Nuber and Diment¹³⁸ successfully treated 2 olecranon stress fractures in competitive pitchers with splinting and cessation of throwing. Both players had radiographic union and returned to pitching.

Operative Treatment

Complete olecranon stress fractures in competitive throwers often require surgical treatment with cannulated screws or plate osteosynthesis. The goal of surgical fixation is compression and rigid fixation across the fracture site. Paci and colleagues¹³⁹ have published the only report on surgical fixation of proximal olecranon stress fractures in baseball players. They performed percutaneous fixation of the proximal ulna stress fracture in 25 baseball players and had follow-up on 18 of the athletes. All 18 fractures went on to union, with 17 of 18 (94%) athletes able to return to preinjury level of play.

OSTEOCHONDRITIS DISSECANS OF THE CAPITELLUM

The radiocapitellar joint experiences compressive forces during valgus stress from overhead throwing motions. Repetitive compressive trauma, in addition to ischemia and genetics, has been

implicated in the formation of OCD of the capitellum. The exact cause, however, remains unclear. A wide spectrum of injuries can result, including subchondral changes to secondary osteochondrosis of the radial head to loose bodies. Treatment depends on the severity and stability of the osteochondral lesion. Patients often complain of lateral elbow pain on palpation and valgus stress. There is frequently an associated loss of elbow extension, ranging from 5° to 20°. Athletes may also have swelling or effusion, tenderness over the lateral aspect of the elbow, and crepitance with motion.¹⁴⁰ Standard radiographs demonstrate classic findings of a subchondral cyst of the capitellum in early cases. In more advanced stages, flattening and irregularity of the capitellar articular surface may be seen.

Mihata and colleagues¹⁴¹ performed a biomechanical cadaveric study and found that OCDs of the radiocapitellar joint increase elbow valgus laxity and contact pressure without increasing UCL strain. Natural history suggests that as many as 50% of patients experience some degeneration in the radiocapitellar joint.¹⁴²

Nonoperative Treatment

Nonoperative management should be reserved for early lesions and consists of a minimum 6-week period of rest from throwing and valgus stress.¹⁴³ After pain resolves, a strengthening program is initiated with isometric exercises, followed by isotonic exercises. Aggressive high-speed, eccentric, and plyometric exercises are progressively included to prepare athletes for the start of an interval throwing program. Mihara and colleagues¹⁴⁴ reported on 39 baseball players with a mean age of 12.8 years with OCD lesions of the capitellum treated nonoperatively. At a mean follow-up of 14.4 months, 25 of 30 early lesions were healed compared with 1 of 9 advanced lesions. A majority of the lesions in patients with open physes healed (16 of 17) compared with only 11 of 22 patients with closed physes.

Operative Treatment

Surgical intervention is limited to patients who do not respond to a nonoperative course of treatment or those with advanced disease resulting in the development of loose fragments within the joint. Initially, surgical management consisted of open débridement and fragment excision. Bauer and colleagues¹⁴⁵ reported long-term outcomes in 31 patients with a mean age of 20 years. Of these patients, 22 had loose body excision and 1 had radial head excision, and most cases seemed to be advanced lesions (20 of 31). At average 23-year

follow-up, there was a 40% recurrence of symptoms and loss of elbow extension, with greater than 60% of examined radiocapitellar joints demonstrating degenerative joint disease. In an initial study, Takahara and colleagues¹⁴² reported 39 patients with an average age of 17.6 years treated with open fragment excision. At average 14.7 years' follow-up, 26% of patients reported good results, and 49% returned to full sports participation. Takahara and colleagues¹⁴⁶ followed their initial study with another study adding 16 patients to their original cohort. The investigators added that better results in terms of pain and radiographic findings were seen in patients with lesions measuring less than 50% of the capitellar articular width.

Arthroscopic débridement and abrasion chondroplasty have had promising results in numerous short and midterm follow-up studies, with pain relief and objective improvements in elbow ROM.^{140,147–150} Byrd and Jones¹⁴⁸ examined 10 patients ages 11 to 16 years, 7 with advanced disease (grade IV or V lesions based on the American Sports Medicine Institute classification system of OCD lesions). At 4-year follow-up, although all 7 lesions healed, only 4 of 10 patients had returned to sport, and 2 demonstrated degenerative radiographic findings. The investigators found no correlation between grade of lesion and postoperative outcomes or return to sport. Ruch and colleagues¹⁵¹ treated 12 patients, ages 11 to 17 years, with unstable elbow lesions (mean size 2.5 cm) for a mean of 3.2 years. Improved extension was seen postoperatively, mechanical symptoms resolved in 11 patients, and 11 of 12 patients were highly satisfied. Radiographs demonstrated capitellar remodeling in all elbows. Only 3 patients, however, returned to sport.

Fragment fixation has been attempted through a variety of techniques and implants.^{146,152–155} Reliable results have been seen in advanced cases, but published data have been limited to small case series. Lateral closing wedge osteotomy has been used to unload the radiocapitellar joint. The procedure is technically challenging because ulnohumeral joint alignment must be maintained. Because the lesion is not directly addressed, this procedure is reserved for early lesions that are stable. Kiyoshige and colleagues¹⁵⁶ evaluated 7 baseball players ages 11 to 18 who underwent a lateral closing wedge osteotomy. At 7- to 12-year follow-up, 6 of 7 patients (86%) had complete relief of pain with return to sport.

Osteochondral autograft transplantation (OAT) is another technique to treat advanced OCD lesions of the capitellum. Contraindications include degenerative changes in the radiocapitellar

compartment and radial head and capitellar deformity. Placement of grafts in the tight lateral compartment make congruent placement of the graft difficult.¹⁵⁷ Early results demonstrate reasonable outcomes with OAT but long-term follow-up in the throwing population is needed before definitive conclusions can be drawn.^{158,159}

SUMMARY

Overhead throwing activities expose the elbow to tremendous valgus stress, making athletes vulnerable to a specific constellation of injuries. Although baseball players, in particular pitchers, are the athletes most commonly affected, overhead athletes in football, volleyball, tennis, and javelin throwing also are affected.

Increasing participation in overhead throwing sports has led to a sharp increase in injuries. Understanding the anatomy and function of the elbow, along with the biomechanical relationship between the two, remains vital to appropriate management. Advances in surgical technique to reconstruct the UCL have led to improved outcomes while multiple fixation devices and grafts have been evaluated.

REFERENCES

1. Nassab PF, Schickendantz MS. Evaluation and treatment of medial ulnar collateral ligament injuries in the throwing athlete. *Sports Med Arthrosc* 2006;14(4):221–31.
2. National Federation of State High School Associations. National Federation of State High School Associations 2012–2013 High school athletics participation survey. 2013.
3. Miller CD, Savoie FH 3rd. Valgus extension injuries of the elbow in the throwing athlete. *J Am Acad Orthop Surg* 1994;2(5):261–9.
4. Morrey BF, Tanaka S, An KN. Valgus stability of the elbow. A definition of primary and secondary constraints. *Clin Orthop Relat Res* 1991;(265):187–95.
5. Schwab GH, Bennett JB, Woods GW, et al. Biomechanics of elbow instability: the role of the medial collateral ligament. *Clin Orthop Relat Res* 1980;(146):42–52.
6. Morrey BF. Applied anatomy and biomechanics of the elbow joint. *Instr Course Lect* 1986;35:59–68.
7. Jobe FW, Kvitne RS. Elbow instability in the athlete. *Instr Course Lect* 1991;40:17–23.
8. Sojbjerg JO, Ovesen J, Nielsen S. Experimental elbow instability after transection of the medial collateral ligament. *Clin Orthop Relat Res* 1987;(218):186–90.
9. Regan WD, Korinek SL, Morrey BF, et al. Biomechanical study of ligaments around the elbow joint. *Clin Orthop Relat Res* 1991;(271):170–9.

10. Callaway GH, Field LD, Deng XH, et al. Biomechanical evaluation of the medial collateral ligament of the elbow. *J Bone Joint Surg Am* 1997; 79(8):1223–31.
11. Davidson PA, Pink M, Perry J, et al. Functional anatomy of the flexor pronator muscle group in relation to the medial collateral ligament of the elbow. *Am J Sports Med* 1995;23(2):245–50.
12. Jobe FW, Moynes DR, Tibone JE, et al. An EMG analysis of the shoulder in pitching. A second report. *Am J Sports Med* 1984;12(3):218–20.
13. Hamilton CD, Glousman RE, Jobe FW, et al. Dynamic stability of the elbow: electromyographic analysis of the flexor pronator group and the extensor group in pitchers with valgus instability. *J Shoulder Elbow Surg* 1996;5(5):347–54.
14. Glousman RE, Barron J, Jobe FW, et al. An electromyographic analysis of the elbow in normal and injured pitchers with medial collateral ligament insufficiency. *Am J Sports Med* 1992;20(3):311–7.
15. Digiovine NM, Jobe FW, Pink M, et al. An electromyographic analysis of the upper extremity in pitching. *J Shoulder Elbow Surg* 1992;1(1):15–25.
16. Sisto DJ, Jobe FW, Moynes DR, et al. An electromyographic analysis of the elbow in pitching. *Am J Sports Med* 1987;15(3):260–3.
17. MacWilliams BA, Choi T, Perezous MK, et al. Characteristic ground-reaction forces in baseball pitching. *Am J Sports Med* 1998;26(1):66–71.
18. Watkins RG, Dennis S, Dillin WH, et al. Dynamic EMG analysis of torque transfer in professional baseball pitchers. *Spine* 1989;14(4):404–8.
19. Pappas AM, Zawacki RM, Sullivan TJ. Biomechanics of baseball pitching. A preliminary report. *Am J Sports Med* 1985;13(4):216–22.
20. Fleisig GS, Andrews JR, Dillman CJ, et al. Kinetics of baseball pitching with implications about injury mechanisms. *Am J Sports Med* 1995;23(2): 233–9.
21. Dines JS, Frank JB, Akerman M, et al. Glenohumeral internal rotation deficits in baseball players with ulnar collateral ligament insufficiency. *Am J Sports Med* 2009;37(3):566–70.
22. Garrison JC, Cole MA, Conway JE, et al. Shoulder range of motion deficits in baseball players with an ulnar collateral ligament tear. *Am J Sports Med* 2012;40(11):2597–603.
23. Meister K, Day T, Horodyski M, et al. Rotational motion changes in the glenohumeral joint of the adolescent/Little League baseball player. *Am J Sports Med* 2005;33(5):693–8.
24. Crockett HC, Gross LB, Wilk KE, et al. Osseous adaptation and range of motion at the glenohumeral joint in professional baseball pitchers. *Am J Sports Med* 2002;30(1):20–6.
25. Borsa PA, Wilk KE, Jacobson JA, et al. Correlation of range of motion and glenohumeral translation in professional baseball pitchers. *Am J Sports Med* 2005;33(9):1392–9.
26. Mihata T, Lee Y, McGarry MH, et al. Excessive humeral external rotation results in increased shoulder laxity. *Am J Sports Med* 2004;32(5):1278–85.
27. Polster JM, Bullen J, Obuchowski NA, et al. Relationship between humeral torsion and injury in professional baseball pitchers. *Am J Sports Med* 2013;41(9):2015–21.
28. Burkhart SS, Morgan CD, Kibler WB. The disabled throwing shoulder: spectrum of pathology Part I: pathoanatomy and biomechanics. *Arthroscopy* 2003;19(4):404–20.
29. Hang DW, Chao CM, Hang YS. A clinical and roentgenographic study of Little League elbow. *Am J Sports Med* 2004;32(1):79–84.
30. Limpisvasti O, ElAttrache NS, Jobe FW. Understanding shoulder and elbow injuries in baseball. *J Am Acad Orthop Surg* 2007;15(3):139–47.
31. Kooima CL, Anderson K, Craig JV, et al. Evidence of subclinical medial collateral ligament injury and posteromedial impingement in professional baseball players. *Am J Sports Med* 2004;32(7):1602–6.
32. Nazarian LN, McShane JM, Ciccotti MG, et al. Dynamic US of the anterior band of the ulnar collateral ligament of the elbow in asymptomatic major league baseball pitchers. *Radiology* 2003;227(1): 149–54.
33. Ellenbecker TS, Mattalino AJ, Elam EA, et al. Medial elbow joint laxity in professional baseball pitchers. A bilateral comparison using stress radiography. *Am J Sports Med* 1998;26(3):420–4.
34. Mair SD, Uhl TL, Robbe RG, et al. Physeal changes and range-of-motion differences in the dominant shoulders of skeletally immature baseball players. *J Shoulder Elbow Surg* 2004;13(5):487–91.
35. King JW, Brelsford HJ, Tullos HS. Analysis of the pitching arm of the professional baseball pitcher. *Clin Orthop Relat Res* 1969;67:116–23.
36. Wilson FD, Andrews JR, Blackburn TA, et al. Valgus extension overload in the pitching elbow. *Am J Sports Med* 1983;11(2):83–8.
37. Yocum LA. The diagnosis and nonoperative treatment of elbow problems in the athlete. *Clin Sports Med* 1989;8(3):439–51.
38. Conway JE, Jobe FW, Glousman RE, et al. Medial instability of the elbow in throwing athletes. Treatment by repair or reconstruction of the ulnar collateral ligament. *J Bone Joint Surg Am* 1992;74(1):67–83.
39. Beals RK. The normal carrying angle of the elbow. A radiographic study of 422 patients. *Clin Orthop Relat Res* 1976;(119):194–6.
40. Childress HM. Recurrent ulnar-nerve dislocation at the elbow. *Clin Orthop Relat Res* 1975;(108):168–73.
41. Del Pizzo W, Jobe FW, Norwood L. Ulnar nerve entrapment syndrome in baseball players. *Am J Sports Med* 1977;5(5):182–5.

42. Morrey BF, An KN. Functional anatomy of the ligaments of the elbow. *Clin Orthop Relat Res* 1985;(201):84–90.
43. Morrey BF, An KN. Articular and ligamentous contributions to the stability of the elbow joint. *Am J Sports Med* 1983;11(5):315–9.
44. Boatright JR, D'Alessandro DF. Nerve entrapment syndromes at the elbow. Operative techniques in upper extremity sports injuries. St Louis (MO): Mosby-Year; 1996. p. 518–37.
45. Azar FM, Andrews JR, Wilk KE, et al. Operative treatment of ulnar collateral ligament injuries of the elbow in athletes. *Am J Sports Med* 2000; 28(1):16–23.
46. Greaney RB, Gerber FH, Laughlin RL, et al. Distribution and natural history of stress fractures in U.S. Marine recruits. *Radiology* 1983;146(2):339–46.
47. Nielsen MB, Hansen K, Holmer P, et al. Tibial periosteal reactions in soldiers. A scintigraphic study of 29 cases of lower leg pain. *Acta Orthop Scand* 1991;62(6):531–4.
48. Zwas ST, Elkanovitch R, Frank G. Interpretation and classification of bone scintigraphic findings in stress fractures. *J Nucl Med* 1987;28(4):452–7.
49. Wilcox JR Jr, Moniot AL, Green JP. Bone scanning in the evaluation of exercise-related stress injuries. *Radiology* 1977;123(3):699–703.
50. Geslien GE, Thrall JH, Espinosa JL, et al. Early detection of stress fractures using 99mTc-polyphosphate. *Radiology* 1976;121(3):683–7.
51. Prather JL, Nusynowitz ML, Snowdy HA, et al. Scintigraphic findings in stress fractures. *J Bone Joint Surg Am* 1977;59(7):869–74.
52. Ammann W, Matheson GO. Radionuclide bone imaging in the detection of stress fractures. *Clin J Sport Med* 1991;1(2):115–22.
53. Roub LW, Gumerman LW, Hanley EN Jr, et al. Bone stress: a radionuclide imaging perspective. *Radiology* 1979;132(2):431–8.
54. Matheson GO, Clement DB, McKenzie DC, et al. Stress fractures in athletes. A study of 320 cases. *Am J Sports Med* 1987;15(1):46–58.
55. Nakanishi K, Masatomi T, Ochi T, et al. MR arthrography of elbow: evaluation of the ulnar collateral ligament of elbow. *Skeletal Radiol* 1996;25(7):629–34.
56. Schwartz ML, al-Zahrani S, Morwessel RM, et al. Ulnar collateral ligament injury in the throwing athlete: evaluation with saline-enhanced MR arthrography. *Radiology* 1995;197(1):297–9.
57. Gaary EA, Potter HG, Altchek DW. Medial elbow pain in the throwing athlete: MR imaging evaluation. *AJR Am J Roentgenol* 1997;168(3):795–800.
58. Potter HG. Imaging of posttraumatic and soft tissue dysfunction of the elbow. *Clin Orthop Relat Res* 2000;(370):9–18.
59. Timmerman LA, Schwartz ML, Andrews JR. Preoperative evaluation of the ulnar collateral ligament by magnetic resonance imaging and computed tomography arthrography. Evaluation in 25 baseball players with surgical confirmation. *Am J Sports Med* 1994;22(1):26–31 [discussion: 32].
60. Kenter K, Behr CT, Warren RF, et al. Acute elbow injuries in the National Football League. *J Shoulder Elbow Surg* 2000;9(1):1–5.
61. Arendt EA, editor. Orthopaedic knowledge update sports medicine. Rosemont (IL): AAOS; 1999.
62. Rettig AC, Sherrill C, Snead DS, et al. Nonoperative treatment of ulnar collateral ligament injuries in throwing athletes. *Am J Sports Med* 2001;29(1): 15–7.
63. Podesta L, Crow SA, Volkmer D, et al. Treatment of partial ulnar collateral ligament tears in the elbow with platelet-rich plasma. *Am J Sports Med* 2013; 41(7):1689–94.
64. Jobe FW, Stark H, Lombardo SJ. Reconstruction of the ulnar collateral ligament in athletes. *J Bone Joint Surg Am* 1986;68(8):1158–63.
65. Norwood LA, Shook JA, Andrews JR. Acute medial elbow ruptures. *Am J Sports Med* 1981;9(1):16–9.
66. Morrey BF, Regan WD. Throwing injuries. In: DeLee JC, Drez D Jr, editors. Orthopaedic sports medicine: principles and practice. Philadelphia: WB Saunders; 1994. p. 882–9.
67. Cain EL Jr, Andrews JR, Dugas JR, et al. Outcome of ulnar collateral ligament reconstruction of the elbow in 1281 athletes: results in 743 athletes with minimum 2-year follow-up. *Am J Sports Med* 2010;38(12):2426–34.
68. Savoie FH 3rd, Trenhaile SW, Roberts J, et al. Primary repair of ulnar collateral ligament injuries of the elbow in young athletes: a case series of injuries to the proximal and distal ends of the ligament. *Am J Sports Med* 2008;36(6):1066–72.
69. Andrews JR, Timmerman LA. Outcome of elbow surgery in professional baseball players. *Am J Sports Med* 1995;23(4):407–13.
70. Savoie FH 3rd, Morgan C, Yaste J, et al. Medial ulnar collateral ligament reconstruction using hamstring allograft in overhead throwing athletes. *J Bone Joint Surg Am* 2013;95(12):1062–6.
71. Smith GR, Altchek DW, Pagnani MJ, et al. A muscle-splitting approach to the ulnar collateral ligament of the elbow. Neuroanatomy and operative technique. *Am J Sports Med* 1996;24(5): 575–80.
72. Thompson WH, Jobe FW, Yocum LA, et al. Ulnar collateral ligament reconstruction in athletes: muscle-splitting approach without transposition of the ulnar nerve. *J Shoulder Elbow Surg* 2001; 10(2):152–7.
73. Langer P, Fadale P, Hulstyn M. Evolution of the treatment options of ulnar collateral ligament injuries of the elbow. *Br J Sports Med* 2006;40(6): 499–506.

74. Armstrong AD, Dunning CE, Ferreira LM, et al. A biomechanical comparison of four reconstruction techniques for the medial collateral ligament-deficient elbow. *J Shoulder Elbow Surg* 2005; 14(2):207–15.
75. Ahmad CS, Lee TQ, ElAttrache NS. Biomechanical evaluation of a new ulnar collateral ligament reconstruction technique with interference screw fixation. *Am J Sports Med* 2003;31(3):332–7.
76. Ciccotti MG, Siegler S, Kuri JA 2nd, et al. Murphy DJt. Comparison of the biomechanical profile of the intact ulnar collateral ligament with the modified Jobe and the Docking reconstructed elbow: an in vitro study. *Am J Sports Med* 2009;37(5):974–81.
77. Dodson CC, Thomas A, Dines JS, et al. Medial ulnar collateral ligament reconstruction of the elbow in throwing athletes. *Am J Sports Med* 2006; 34(12):1926–32.
78. Hurbanek JG, Anderson K, Crabtree S, et al. Biomechanical comparison of the docking technique with and without humeral bioabsorbable interference screw fixation. *Am J Sports Med* 2009;37(3):526–33.
79. Large TM, Coley ER, Peindl RD, et al. A biomechanical comparison of 2 ulnar collateral ligament reconstruction techniques. *Arthroscopy* 2007;23(2):141–50.
80. McAdams TR, Lee AT, Centeno J, et al. Two ulnar collateral ligament reconstruction methods: the docking technique versus bioabsorbable interference screw fixation—a biomechanical evaluation with cyclic loading. *J Shoulder Elbow Surg* 2007; 16(2):224–8.
81. Morgan RJ, Starman JS, Habet NA, et al. A biomechanical evaluation of ulnar collateral ligament reconstruction using a novel technique for ulnar-sided fixation. *Am J Sports Med* 2010;38(7): 1448–55.
82. Nissen CW. Effectiveness of interference screw fixation in ulnar collateral ligament reconstruction. *Orthopedics* 2008;31(7):646.
83. Paletta GA Jr, Klepps SJ, Difelice GS, et al. Biomechanical evaluation of 2 techniques for ulnar collateral ligament reconstruction of the elbow. *Am J Sports Med* 2006;34(10):1599–603.
84. Paletta GA Jr, Wright RW. The modified docking procedure for elbow ulnar collateral ligament reconstruction: 2-year follow-up in elite throwers. *Am J Sports Med* 2006;34(10):1594–8.
85. Shah RP, Lindsey DP, Sungar GW, et al. An analysis of four ulnar collateral ligament reconstruction procedures with cyclic valgus loading. *J Shoulder Elbow Surg* 2009;18(1):58–63.
86. Rohrbough JT, Altchek DW, Hyman J, et al. Medial collateral ligament reconstruction of the elbow using the docking technique. *Am J Sports Med* 2002;30(4):541–8.
87. Starman JS, Morgan RJ, Fleischli JE, et al. Ulnar collateral ligament reconstruction using the Toggle-Loc with ZipLoop for ulnar side fixation. *Orthopedics* 2010;33(5):312–6.
88. Bowers AL, Dines JS, Dines DM, et al. Elbow medial ulnar collateral ligament reconstruction: clinical relevance and the docking technique. *J Shoulder Elbow Surg* 2010;19(Suppl 2):110–7.
89. Koh JL, Schafer MF, Keuter G, et al. Ulnar collateral ligament reconstruction in elite throwing athletes. *Arthroscopy* 2006;22(11):1187–91.
90. Watson JN, McQueen P, Hutchinson MR. A systematic review of ulnar collateral ligament reconstruction techniques. *Am J Sports Med* 2013. [Epub ahead of print].
91. Dines JS, ElAttrache NS, Conway JE, et al. Clinical outcomes of the DANE TJ technique to treat ulnar collateral ligament insufficiency of the elbow. *Am J Sports Med* 2007;35(12):2039–44.
92. Domb BG, Davis JT, Alberta FG, et al. Clinical follow-up of professional baseball players undergoing ulnar collateral ligament reconstruction using the new Kerlan-Jobe Orthopaedic Clinic overhead athlete shoulder and elbow score (KJOC Score). *Am J Sports Med* 2010;38(8):1558–63.
93. Vitale MA, Ahmad CS. The outcome of elbow ulnar collateral ligament reconstruction in overhead athletes: a systematic review. *Am J Sports Med* 2008;36(6):1193–205.
94. Petty DH, Andrews JR, Fleisig GS, et al. Ulnar collateral ligament reconstruction in high school baseball players: clinical results and injury risk factors. *Am J Sports Med* 2004;32(5):1158–64.
95. Lee GH, Limpisvasti O, Park MC, et al. Revision ulnar collateral ligament reconstruction using a suspension button fixation technique. *Am J Sports Med* 2010;38(3):575–80.
96. Dines JS, Yocum LA, Frank JB, et al. Revision surgery for failed elbow medial collateral ligament reconstruction. *Am J Sports Med* 2008;36(6): 1061–5.
97. Schickendantz MS. Diagnosis and treatment of elbow disorders in the overhead athlete. *Hand Clin* 2002;18(1):65–75.
98. Li X, Dines JS, Gorman M, et al. Anconeus epitrochlearis as a source of medial elbow pain in baseball pitchers. *Orthopedics* 2012;35(7): e1129–32.
99. Aoki M, Kanaya K, Aiki H, et al. Cubital tunnel syndrome in adolescent baseball players: a report of six cases with 3- to 5-year follow-up. *Arthroscopy* 2005;21(6):758.
100. Bradshaw DY, Shefner JM. Ulnar neuropathy at the elbow. *Neurol Clin* 1999;17(3):447–61, v–vi.
101. Huang JH, Samadani U, Zager EL. Ulnar nerve entrapment neuropathy at the elbow: simple decompression. *Neurosurgery* 2004;55(5):1150–3.

102. Rokito AS, Iviciviahon PJ, Jobe FW. Cubital tunnel syndrome. *Oper Tech Sports Med* 1996;4(1): 15–20.
103. Glousman RE. Ulnar nerve problems in the athlete's elbow. *Clin Sports Med* 1990;9(2):365–77.
104. Gabel GT, Morrey BF. Operative treatment of medical epicondylitis. Influence of concomitant ulnar neuropathy at the elbow. *J Bone Joint Surg Am* 1995;77(7):1065–9.
105. Griffin LY. Orthopaedic knowledge update sports medicine. Rosemont (IL): AAOS; 1994. p. 179–90.
106. Pechan J, Julis I. The pressure measurement in the ulnar nerve. A contribution to the pathophysiology of the cubital tunnel syndrome. *J Biomech* 1975; 8(1):75–9.
107. Chen FS, Rokito AS, Jobe FW. Medial elbow problems in the overhead-throwing athlete. *J Am Acad Orthop Surg* 2001;9(2):99–113.
108. Wei SH, Jong YJ, Chang YJ. Ulnar nerve conduction velocity in injured baseball pitchers. *Arch Phys Med Rehabil* 2005;86(1):21–5 [quiz: 180].
109. Cain EL Jr, Dugas JR, Wolf RS, et al. Elbow injuries in throwing athletes: a current concepts review. *Am J Sports Med* 2003;31(4):621–35.
110. Rettig AC, Ebben JR. Anterior subcutaneous transfer of the ulnar nerve in the athlete. *Am J Sports Med* 1993;21(6):836–9 [discussion: 839–40].
111. Zlowodzki M, Chan S, Bhandari M, et al. Anterior transposition compared with simple decompression for treatment of cubital tunnel syndrome. A meta-analysis of randomized, controlled trials. *J Bone Joint Surg Am* 2007;89(12):2591–8.
112. Macadam SA, Gandhi R, Bezuhly M, et al. Simple decompression versus anterior subcutaneous and submuscular transposition of the ulnar nerve for cubital tunnel syndrome: a meta-analysis. *J Hand Surg* 2008;33(8):1314.e1–2.
113. Larson RL, Singer KM, Bergstrom R, et al. Little league survey: the Eugene study. *Am J Sports Med* 1976;4(5):201–9.
114. Gugenheim JJ Jr, Stanley RF, Woods GW, et al. Little League survey: the Houston study. *Am J Sports Med* 1976;4(5):189–200.
115. Grana WA, Rashkin A. Pitcher's elbow in adolescents. *Am J Sports Med* 1980;8(5):333–6.
116. Benjamin HJ, Briner WW Jr. Little league elbow. *Clin J Sport Med* 2005;15(1):37–40.
117. Lyman S, Fleisig GS, Andrews JR, et al. Effect of pitch type, pitch count, and pitching mechanics on risk of elbow and shoulder pain in youth baseball pitchers. *Am J Sports Med* 2002;30(4):463–8.
118. Lyman S, Fleisig GS, Waterbor JW, et al. Longitudinal study of elbow and shoulder pain in youth baseball pitchers. *Med Sci Sports Exerc* 2001; 33(11):1803–10.
119. Fleisig GS, Barrentine SW, Zheng N, et al. Kinematic and kinetic comparison of baseball pitching among various levels of development. *J Biomech* 1999;32(12):1371–5.
120. Bennett GE. Elbow and shoulder lesions of baseball players. *Am J Surg* 1959;98:484–92.
121. Wei AS, Khana S, Limpisvasti O, et al. Clinical and magnetic resonance imaging findings associated with Little League elbow. *J Pediatr Orthop* 2010; 30(7):715–9.
122. Harada M, Takahara M, Hirayama T, et al. Outcome of nonoperative treatment for humeral medial epicondylar fragmentation before epiphyseal closure in young baseball players. *Am J Sports Med* 2012;40(7):1583–90.
123. Torg JS. The little league pitcher. *Am Fam Physician* 1972;6(2):71–6.
124. Woods GW, Tullos HS. Elbow instability and medial epicondyle fractures. *Am J Sports Med* 1977;5(1): 23–30.
125. Hines RF, Herndon WA, Evans JP. Operative treatment of medial epicondyle fractures in children. *Clin Orthop Relat Res* 1987;(223):170–4.
126. Ireland ML, Andrews JR. Shoulder and elbow injuries in the young athlete. *Clin Sports Med* 1988; 7(3):473–94.
127. Lawrence JT, Patel NM, Macknlin J, et al. Return to competitive sports after medial epicondyle fractures in adolescent athletes: results of operative and nonoperative treatment. *Am J Sports Med* 2013;41(5):1152–7.
128. Loftice J, Fleisig GS, Zheng N, et al. Biomechanics of the elbow in sports. *Clin Sports Med* 2004;23(4): 519–30, vii–viii.
129. Cohen SB, Valko C, Zoga A, et al. Posteromedial elbow impingement: magnetic resonance imaging findings in overhead throwing athletes and results of arthroscopic treatment. *Arthroscopy* 2011; 27(10):1364–70.
130. Wilk KE, Macrina LC, Cain EL, et al. Rehabilitation of the Overhead Athlete's Elbow. *Sports Health* 2012;4(5):404–14.
131. Andrews JR, St Pierre RK, Carson WG Jr. Arthroscopy of the elbow. *Clin Sports Med* 1986;5(4): 653–62.
132. Timmerman LA, Andrews JR. Undersurface tear of the ulnar collateral ligament in baseball players. A newly recognized lesion. *Am J Sports Med* 1994; 22(1):33–6.
133. Reddy AS, Kvitne RS, Yocum LA, et al. Arthroscopy of the elbow: a long-term clinical review. *Arthroscopy* 2000;16(6):588–94.
134. Kamineni S, ElAttrache NS, O'Driscoll SW, et al. Medial collateral ligament strain with partial posteromedial olecranon resection. A biomechanical study. *J Bone Joint Surg Am* 2004;86A(11): 2424–30.
135. Ahmad CS, Park MC, Elattrache NS. Elbow medial ulnar collateral ligament insufficiency alters

- posteromedial olecranon contact. *Am J Sports Med* 2004;32(7):1607–12.
136. Griggs SM, Weiss AP. Bony injuries of the wrist, forearm, and elbow. *Clin Sports Med* 1996;15(2):373–400.
 137. Schickendantz MS, Ho CP, Koh J. Stress injury of the proximal ulna in professional baseball players. *Am J Sports Med* 2002;30(5):737–41.
 138. Nuber GW, Diment MT. Olecranon stress fractures in throwers. A report of two cases and a review of the literature. *Clin Orthop Relat Res* 1992;(278):58–61.
 139. Paci JM, Dugas JR, Guy JA, et al. Cannulated screw fixation of refractory olecranon stress fractures with and without associated injuries allows a return to baseball. *Am J Sports Med* 2013;41(2):306–12.
 140. Ruchelsman DE, Hall MP, Youm T. Osteochondritis dissecans of the capitellum: current concepts. *J Am Acad Orthop Surg* 2010;18(9):557–67.
 141. Mihata T, Quigley R, Robicheaux G, et al. Biomechanical characteristics of osteochondral defects of the humeral capitellum. *Am J Sports Med* 2013;41(8):1909–14.
 142. Takahara M, Ogino T, Sasaki I, et al. Long term outcome of osteochondritis dissecans of the humeral capitellum. *Clin Orthop Relat Res* 1999;(363):108–15.
 143. Takahara M, Ogino T, Fukushima S, et al. Nonoperative treatment of osteochondritis dissecans of the humeral capitellum. *Am J Sports Med* 1999;27(6):728–32.
 144. Mihara K, Tsutsui H, Nishinaka N, et al. Nonoperative treatment for osteochondritis dissecans of the capitellum. *Am J Sports Med* 2009;37(2):298–304.
 145. Bauer M, Jonsson K, Josefsson PO, et al. Osteochondritis dissecans of the elbow: a long-term follow-up study. *Clin Orthop Relat Res* 1992;284:156–60.
 146. Takahara M, Mura N, Sasaki J, et al. Classification, treatment, and outcome of osteochondritis dissecans of the humeral capitellum. *J Bone Joint Surg Am* 2007;89(6):1205–14.
 147. Baumgarten TE, Andrews JR, Satterwhite YE. The arthroscopic classification and treatment of osteochondritis dissecans of the capitellum. *Am J Sports Med* 1998;26(4):520–3.
 148. Byrd JW, Jones KS. Arthroscopic surgery for isolated capitellar osteochondritis dissecans in adolescent baseball players: minimum three-year follow-up. *Am J Sports Med* 2002;30(4):474–8.
 149. Brownlow HC, O'Connor-Read LM, Perko M. Arthroscopic treatment of osteochondritis dissecans of the capitellum. *Knee Surg Sports Traumatol Arthrosc* 2006;14(2):198–202.
 150. Rahusen FT, Brinkman JM, Eygendaal D. Results of arthroscopic debridement for osteochondritis dissecans of the elbow. *Br J Sports Med* 2006;40(12):966–9.
 151. Ruch DS, Cory JW, Poehling GG. The arthroscopic management of osteochondritis dissecans of the adolescent elbow. *Arthroscopy* 1998;14(8):797–803.
 152. Yadao MA, Field LD, Savoie FH 3rd. Osteochondritis dissecans of the elbow. *Instr Course Lect* 2004;53:599–606.
 153. Harada M, Ogino T, Takahara M, et al. Fragment fixation with a bone graft and dynamic staples for osteochondritis dissecans of the humeral capitellum. *J Shoulder Elbow Surg* 2002;11(4):368–72.
 154. Takeda H, Watarai K, Matsushita T, et al. A surgical treatment for unstable osteochondritis dissecans lesions of the humeral capitellum in adolescent baseball players. *Am J Sports Med* 2002;30(5):713–7.
 155. Kuwahata Y, Inoue G. Osteochondritis dissecans of the elbow managed by Herbert screw fixation. *Orthopedics* 1998;21(4):449–51.
 156. Kiyoshige Y, Takagi M, Yuasa K, et al. Closed-Wedge osteotomy for osteochondritis dissecans of the capitellum. A 7- to 12-year follow-up. *Am J Sports Med* 2000;28(4):534–7.
 157. Miyamoto W, Yamamoto S, Kii R, et al. Oblique osteochondral plugs transplantation technique for osteochondritis dissecans of the elbow joint. *Knee Surg Sports Traumatol Arthrosc* 2009;17(2):204–8.
 158. Iwasaki N, Kato H, Ishikawa J, et al. Autologous osteochondral mosaicplasty for osteochondritis dissecans of the elbow in teenage athletes. *J Bone Joint Surg Am* 2009;91(10):2359–66.
 159. Shimada K, Yoshida T, Nakata K, et al. Reconstruction with an osteochondral autograft for advanced osteochondritis dissecans of the elbow. *Clin Orthop Relat Res* 2005;(435):140–7.
 160. Farrow LD, Mahoney AJ, Stefancin JJ, et al. Quantitative analysis of the medial ulnar collateral ligament ulnar footprint and its relationship to the ulnar sublime tubercle. *Am J Sports Med* 2011;39(9):1936–41.